

Current Opioid Misuse Measure (COMM)[™]

The Current Opioid Misuse Measure (COMM)[™] is a brief patient self-assessment to monitor chronic pain patients on opioid therapy. The COMM[™] was developed with guidance from a group of pain and addiction experts and input from pain management clinicians in the field. Experts and providers identified six key issues to determine if patients already on long-term opioid treatment are exhibiting aberrant medication-related behaviors:

- *Signs & Symptoms of Intoxication*
- *Emotional Volatility*
- *Evidence of Poor Response to Medications*
- *Addiction*
- *Healthcare Use Patterns*
- *Problematic Medication Behavior*

The COMM[™] will help clinicians identify whether a patient, currently on long-term opioid therapy, may be exhibiting aberrant behaviors associated with misuse of opioid medications. In contrast, the Screener and Opioid Assessment for Patients with Pain (SOAPP®) is intended to predict which patients, being considered for long-term opioid therapy, may exhibit aberrant medications behaviors in the future. Since the COMM[™] examines concurrent misuse, it is ideal for helping clinicians monitor patients' aberrant medication-related behaviors over the course of treatment. The COMM[™] is:

- A quick and easy to administer patient-self assessment
- 17 items
- Simple to score
- Completed in less than 10 minutes
- Validated with a group of approximately 500 chronic pain patients on opioid therapy
- Ideal for documenting decisions about the level of monitoring planned for a particular patient or justifying referrals to specialty pain clinic.
- The COMM[™] is for clinician use only. The tool is not meant for commercial distribution.
- The COMM[™] is **NOT** a lie detector. Patients determined to misrepresent themselves will still do so. Other clinical information should be used with COMM[™] scores to decide if and when modifications to particular patient's treatment plan is needed.
- It is important to remember that all chronic pain patients deserve treatment of their pain. Providers who are not comfortable treating certain patients should refer those patients to a specialist.

©2008 Inflexion, Inc. Permission granted solely for use in published format by individual practitioners in clinical practice. No other uses or alterations are authorized or permitted by copyright holder. Permissions questions: PainEDU@inflexion.com. The COMM[™] was developed with a grant from the National Institutes of Health and an educational grant from Endo Pharmaceuticals.

COMM™

Please answer each question as honestly as possible. Keep in mind that we are only asking about the **past 30 days**. There are no right or wrong answers. If you are unsure about how to answer the question, please give the best answer you can.

Please answer the questions using the following scale:	Never	Seldom	Sometimes	Often	Very Often
	0	1	2	3	4
1. In the past 30 days, how often have you had trouble with thinking clearly or had memory problems?	<input type="radio"/>				
2. In the past 30 days, how often do people complain that you are not completing necessary tasks? (i.e., doing things that need to be done, such as going to class, work or appointments)	<input type="radio"/>				
3. In the past 30 days, how often have you had to go to someone other than your prescribing physician to get sufficient pain relief from medications? (i.e., another doctor, the Emergency Room, friends, street sources)	<input type="radio"/>				
4. In the past 30 days, how often have you taken your medications differently from how they are prescribed?	<input type="radio"/>				
5. In the past 30 days, how often have you seriously thought about hurting yourself?	<input type="radio"/>				
6. In the past 30 days, how much of your time was spent thinking about opioid medications (having enough, taking them, dosing schedule, etc.)?	<input type="radio"/>				

©2008 Inflexxion, Inc. Permission granted solely for use in published format by individual practitioners in clinical practice. No other uses or alterations are authorized or permitted by copyright holder. Permissions questions: PainEDU@inflexxion.com. The COMM™ was developed with a grant from the National Institutes of Health and an educational grant from Endo Pharmaceuticals.

Please answer the questions using the following scale:	Never	Seldom	Sometimes	Often	Very Often
	0	1	2	3	4
7. In the past 30 days, how often have you been in an argument?	<input type="radio"/>				
8. In the past 30 days, how often have you had trouble controlling your anger (e.g., road rage, screaming, etc.)?	<input type="radio"/>				
9. In the past 30 days, how often have you needed to take pain medications belonging to someone else?	<input type="radio"/>				
10. In the past 30 days, how often have you been worried about how you're handling your medications?	<input type="radio"/>				
11. In the past 30 days, how often have others been worried about how you're handling your medications?	<input type="radio"/>				
12. In the past 30 days, how often have you had to make an emergency phone call or show up at the clinic without an appointment?	<input type="radio"/>				
13. In the past 30 days, how often have you gotten angry with people?	<input type="radio"/>				
14. In the past 30 days, how often have you had to take more of your medication than prescribed?	<input type="radio"/>				
15. In the past 30 days, how often have you borrowed pain medication from someone else?	<input type="radio"/>				
16. In the past 30 days, how often have you used your pain medicine for symptoms other than for pain (e.g., to help you sleep, improve your mood, or relieve stress)?	<input type="radio"/>				

©2008 Inflexxion, Inc. Permission granted solely for use in published format by individual practitioners in clinical practice. No other uses or alterations are authorized or permitted by copyright holder. Permissions questions: PainEDU@inflexxion.com. The COMM™ was developed with a grant from the National Institutes of Health and an educational grant from Endo Pharmaceuticals.

Please answer the questions using the following scale:	Never	Seldom	Sometimes	Often	Very Often
	0	1	2	3	4
17. In the past 30 days, how often have you had to visit the Emergency Room?	○	○	○	○	○

©2008 Inflexion, Inc. Permission granted solely for use in published format by individual practitioners in clinical practice. No other uses or alterations are authorized or permitted by copyright holder. Permissions questions: PainEDU@inflexion.com. The COMM™ was developed with a grant from the National Institutes of Health and an educational grant from Endo Pharmaceuticals.



Scoring Instructions for the COMM™

To score the COMM™, simply add the rating of all the questions. A score of 9 or higher is considered a positive

Sum of Questions	COMM Indication
> or = 9	+
< 9	-

As for any scale, the results depend on what cutoff score is chosen. A score that is sensitive in detecting patients who are abusing or misusing their opioid medication will necessarily include a number of patients that are not really abusing or misusing their medication. The COMM™ was intended to over-identify misuse, rather than to mislabel someone as responsible when they are not. This is why a low cut-off score was accepted. We believe that it is more important to identify patients who have only a possibility of misusing their medications than to fail to identify those who are actually abusing their medication. Thus, it is possible that the COMM™ will result in false positives – patients identified as misusing their medication when they were not.

The table below presents several statistics that describe how effective the COMM™ is at different cutoff values. These values suggest that the COMM™ is a sensitive test. This confirms that the COMM™ is better at identifying who is misusing their medication than identifying who is not misusing. Clinically, a score of 9 or higher will identify 77% of those who actually turn out to be at high risk. The Negative Predictive Values for a cutoff score of 9 is .95, which means that most people who have a negative COMM™ are likely not misusing their medication. Finally, the Positive likelihood ratio suggests that a positive COMM™ score (at a cutoff of 9) is nearly 3 times (3.48 times) as likely to come from someone who is actually misusing their medication (note that, of these statistics, the likelihood ratio is least affected by prevalence rates). All this implies that by using a cutoff score of 9 will ensure that the provider is least likely to miss someone who is really misusing their prescription opioids. However, one should remember that a low COMM™ score suggests the patient is really at low-risk, while a high COMM™ score will contain a larger percentage of false positives (about 34%), while at the same time retaining a large percentage of true positives. This could be improved, so that a positive score has a lower false positive rate, but only at the risk of missing more of those who actually do show aberrant behavior.

COMM™ Cutoff Score	Sensitivity	Specificity	Positive Predictive Value	Negative Predictive Value	Positive Likelihood Ratio	Negative Likelihood Ration
Score 9 or above	.77	.66	.66	.95	3.48	.08

©2008 Inflexion, Inc. Permission granted solely for use in published format by individual practitioners in clinical practice. No other uses or alterations are authorized or permitted by copyright holder. Permissions questions: PainEDU@inflexion.com. The COMM™ was developed with a grant from the National Institutes of Health and an educational grant from Endo Pharmaceuticals.